

On principal fibrations associated with one algebra

Maria Trnková^{1,2}

email: Maria-Trnkova@seznam.cz

¹Department of Algebra and Geometry, Faculty of Science,
Palacky University in Olomouc, Tomkova 10, Olomouc, Czech Republic

²Department of Geometry, NIMM,
Kazan State University, prof. Nuzhina 1/37, Kazan, Russia

To the memory of B.N. Shapukov

Abstract *In this paper we study two types of fibrations associated with a 3-dimensional unital associative irreducible algebra and their basic properties. We investigate trivial principal fibrations of degenerate semi-Euclidean sphere and their semi-conformal and projective models. We use Norden normalization method for constructing second model.*

1. PRINCIPAL FIBRATIONS OF G BY ITS SUBGROUPS

Let us denote by $\mathfrak{A}$ a unital associative n -dimensional algebra with multiplication xy , by $G \subset \mathfrak{A}$ the set of invertible elements. It is well known that G is a Lie group with the same multiplication rule. Let $\mathfrak{B}$ be a unital subalgebra of algebra $\mathfrak{A}$ and $H \subset G$ the set of invertible elements. So H is a Lie subgroup of group G . We consider the factor-space G/H of right cosets. A fibration $(G, \pi, M = G/H)$ is a principal fibration with the structure group H , where π is a canonical projection [1]. Here we have a fibration:

$$H \rightarrow G \rightarrow G/H.$$

We consider all 3-dimensional unital associative irreducible algebras up to isomorphism. It is known that there exist only tree types of such algebras [2], [3]. In special choice of basic units they have the following multiplication rules:

I)

	1	e_1	e_2
1	1	e_1	e_2
e_1	e_1	e_2	0
e_2	e_2	0	0

II)

	1	e_1	e_2
1	1	e_1	e_2
e_1	e_1	1	e_2
e_2	e_2	$-e_2$	0

III)

	1	e_1	e_2
1	1	e_1	e_2
e_1	e_1	0	0
e_2	e_2	0	0

For these algebras N.Belova found all subalgebras with unit and principal fibrations [4]. Only in algebras of type II and III exist the associated conjugation $x = x_0 + x^i e_i \rightarrow \bar{x} = x_0 - x^i e_i$ which has the property $\overline{xy} = \bar{y} \bar{x}$.

We consider the bilinear form

$$(x, y) = \frac{1}{2}(x\bar{y} + y\bar{x}). \quad (1)$$

For algebras of type II and III this form takes the real values and it determines a degenerate scalar product on them:

$$\text{II})(x, y) = x_0 y_0 - x_1 y_1, \quad \text{III})(x, y) = x_0 y_0.$$

So these algebras have a structure of semi-Euclidean vector spaces with rank 2 and 1, respectively. In our discussion we concentrate on the algebra of type II, because it is less degenerate than type III.

This is non-Abelian algebra. Multiplication rule for algebra elements is:

$$xy = (x_0 + x_1 e_1 + x_2 e_2)(y_0 + y_1 e_1 + y_2 e_2) = \quad (2)$$

$$x_0 y_0 + x_1 y_1 + (x_0 y_1 + x_1 y_0) e_1 + (x_2(y_0 - y_1) + (x_0 + x_1)y_2) e_2.$$

An inverse element of x is:

$$x^{-1} = \frac{x_0 - x_1 e_1 - x_2 e_2}{(x_0)^2 - (x_1)^2}. \quad (3)$$

The set of invertible elements $G = \{x \in \mathfrak{A} \mid (x_0)^2 - (x_1)^2 \neq 0\}$ is a non-Abelian Lie group. Its underlying manifold is $\mathbb{R}^3$ without two transversal 2-planes, hence it consists from 4 connected components.

Proposition(N.Belova) *Any 2-dimensional subalgebra of algebra of type II with an algebra unit is isomorphic to double or dual numbers algebras.*

We consider a subalgebra $R(e_1)$ with basis $\{1, e_1\}$, it is an algebra of double numbers, and a subalgebra $R(e_2)$ with basis $\{1, e_2\}$, it is an algebra of dual numbers. The set of their invertible elements $H_1 = \{x_0 + x_1 e_1 \in R(e_1) \mid x_0^2 - x_1^2 \neq 0\}$ and $H_2 = \{x_0 + x_2 e_2 \in R(e_2) \mid x_0 \neq 0\}$ are Lie subgroups of the Lie group G . First we take the space of right cosets by $R(e_1)$ subalgebra and fibration by it. Then we have the following proposition:

Proposition (N.Belova) *Fibration $(G, \pi, M = G/H_1)$ determined by formula*

$$\pi(x) = \frac{x_2}{x_0 - x_1} \quad (4)$$

is a trivial principal fibration over the real line $\mathbb{R}$. The typical fiber, it is a plane without two transversal lines. The structure group is H_1 .

Therefore, the manifold of the group G is diffeomorphic to direct sum $\mathbb{R} \times H_1$. The equation of fibers is:

$$u(x_0 - x_1) - x_2 = 0, \quad u \in R. \quad (5)$$

This is 1-parametric family of planes with common axis: $x_0 - x_1 = 0, x_2 = 0$. Of course it is necessary to remove the intersection points of these planes with planes: $x_0 \pm x_1 = 0$.

We consider the left multiplications $x' = ax$ on the group G . They form the 3-parametric Lie group of linear transformations:

$$L(a) = \begin{pmatrix} a_0 & a_1 & 0 \\ a_1 & a_0 & 0 \\ a_2 & -a_2 & a_0 + a_1 \end{pmatrix}, \quad (6)$$

where determinant $\det L(a) = (a_0^2 - a_1^2)(a_0 + a_1) \neq 0$. The multiplications preserve fibers if and only if when $a \in H_1$. This group consists of four connected components.

The right multiplications $x' = xb$ on the group G forms the 3-parametric Lie group of linear transformations:

$$R(b) = \begin{pmatrix} b_0 & b_1 & 0 \\ b_1 & b_0 & 0 \\ b_2 & b_2 & b_0 - b_1 \end{pmatrix}. \quad (7)$$

They preserve the fibration and induce the 2-parametric group of affine transformations on the base:

$$u' = \alpha u + \beta,$$

here $\alpha = \frac{b_0 - b_1}{b_0 + b_1} \neq 0, \beta = \pi(b)$.

The right multiplications include involutions of hyperbolic type with matrices:

$$R(b_2) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ b_2 & b_2 & -1 \end{pmatrix}, \quad R^2 = \text{Id}.$$

They induce the 1-parametric family involutions on the base:

$$u' = -u + b_2.$$

Let us consider rotations and anti-rotations of this semi-Euclidean space. Multiplications $x' = ax$ and $x' = xa$, where $|a|^2 = \pm 1$, are rotations and anti-rotations because of the relations:

$$|x'|^2 = |a|^2 |x|^2 = \pm |x|^2.$$

Each element $a \in \mathfrak{A}$, $|a|^2 = \pm 1$, can be represented as:

$$a = \cosh \varphi + a_0 \sinh \varphi \quad \text{or} \quad a = \sinh \varphi + a_0 \cosh \varphi,$$

where $a_0 = -\bar{a}_0$, $|a_0|^2 = -1$. The bilinear form (1) in the algebra $\mathfrak{A}$ takes real values, therefore it is possible to present it as: $(x, y) = \frac{1}{2}(x\bar{y} + y\bar{x}) = \frac{1}{2}(\bar{x}y + \bar{y}x)$. Consequently, the hyperbolic cosines or sines of angle between x and x' is equal:

$$\frac{(x, ax)}{|x||ax|} = \frac{1/2(x\bar{a}\bar{x} + ax\bar{x})}{|x|^2} = \frac{1/2(x\bar{x}\bar{a} + ax\bar{x})}{|x|^2} = \frac{1}{2}(\bar{a} + a),$$

$$\frac{(x, xa)}{|x||xa|} = \frac{1/2(\bar{x}xa + \bar{x}\bar{a}x)}{|x|^2} = \frac{1/2(\bar{x}xa + \bar{a}\bar{x}x)}{|x|^2} = \frac{1}{2}(a + \bar{a}).$$

In the last two equations we get $\cosh \varphi$ if $|a|^2 = 1$ and we get $\sinh \varphi$ if $|a|^2 = -1$. It means that this angle does not depend on x .

Transformations

$$x' = axb, \quad (8)$$

where $|a|^2 = \pm 1$, $|b|^2 = \pm 1$, are compositions of rotations or anti-rotations $x' = ax$ and $x' = xb$. So, there are proper rotations or anti-rotations.

Transformations

$$x' = a\bar{x}b \quad (9)$$

are compositions of (8) and reflection $x' = \bar{x}$. So, there are improper rotations or anti-rotations.

Lemma *Any proper or improper rotation and anti-rotation of this semi-Euclidean space can be represented by (8) or (9).*

Proof These rotations and anti-rotations are compositions of odd and even numbers of reflections of planes passing through the origin. To each plane corresponds its orthonormal vector n . If vectors x_1 and n are collinear, then $\bar{x}_1 n = \bar{n} x_1$ and $x'_1 = -n\bar{x}_1 n = -n\bar{n} x_1 = -x_1$. If vectors x_2 and n are orthogonal, then $\bar{x}_2 n + \bar{n} x_2 = 0$ and $x'_2 = -n\bar{x}_2 n = n\bar{n} x_2 = x_2$. On the other hand any vector x can be represented as a sum of vectors x_1 and x_2 . It means, that a reflection of plane is: $x' = -n\bar{x}n$. So, composition of odd and even numbers of planes reflections are transformation (8) or (9). $\square$

Let us introduce adapted coordinates (u, λ, φ) to fibration in semi-Euclidean space, here u is a basic coordinate, λ, φ are fiber coordinates. If $|x|^2 > 0$, we denote $\lambda = \pm\sqrt{x_0^2 - x_1^2} \neq 0$, the sign of λ is equal to the sign of x_0 . The adapted coordinates of fibration in this case are:

$$x_0 = \lambda \cosh \varphi, \quad x_1 = \lambda \sinh \varphi, \quad x_2 = u\lambda \exp \varphi, \quad (10)$$

where $\lambda \in \mathbb{R}_0$, $u, \varphi \in \mathbb{R}$.

If $|x|^2 < 0$, then we write $\lambda = \pm\sqrt{x_1^2 - x_0^2}$, the sign of λ is equal to the sign of x_1 :

$$x_0 = \lambda \sinh \varphi, \quad x_1 = \lambda \cosh \varphi, \quad x_2 = u\lambda \exp \varphi. \quad (11)$$

The structure group acts as follows:

$$u' = u, \quad \lambda' = \lambda\rho, \quad \varphi' = \varphi + \psi, \quad (12)$$

where the element $a(0, \rho, \psi)$ of structure group acts on the element $x(u, \lambda, \varphi) \in G$. This group consists of 4 connected components.

2. THE PRINCIPAL SUBFIBRATION OF THE FIBRATION $(G, \pi, M = G/H_1)$

As we already said, the scalar product in the algebra $\mathfrak{A}$ of type II is: $(x, y) = x_0 y_0 - x_1 y_1$. So, the algebra $\mathfrak{A}$ is a 3-dimensional semi-Euclidean space with

rank 2. We call *semi-Euclidean sphere with unit radius* the set of all elements of algebra $\mathfrak{A}$ whose square is equal to one,

$$S^2(1) = \{x \in \mathfrak{A} \mid x_0^2 - x_1^2 = 1\}.$$

It looks like a hyperbolic cylinder. The set of elements with imaginary unit module $|x|^2 = -1$ we call *semi-Euclidean sphere with imaginary unit radius* $S^2(-1)$. One of these spheres can be obtained up from another one by rotation.

We consider the subfibration of the fibration $(G, \pi, M = G/H_1)$ to semi-Euclidean sphere $S^2(1)$, i.e. the fibration $\pi : S^2(1) \rightarrow M$. The fibers of new fibration are intersections of $S^2(1)$ and planes (5). The restriction of the group of double numbers H_1 to $S^2(1)$ is a Lie subgroup S_1 of double numbers with unit module

$$S_1 = \{a_0 + a_1 e_1 \in H_1 \mid a_0^2 - a_1^2 = 1\}.$$

This group consists of two connected components.

Proposition *The fibration $(S^2(1), \pi, M)$ is principal fibration of the group $S^2(1)$ by the Lie subgroup S_1 to right cosets.*

Proof. Let x and y be two sphere points from one fiber. They are also from the same fiber of the principal fibration (G, π, M) . So, there exists a unique element $a \in H_1$ such that $y = ax$. Then $|y|^2 = |a|^2|x|^2$ and hence $|a|^2 = 1$. It means that $a \in S_1$. $\square$

We define coordinates adapted to the fibration on semi-Euclidean sphere $S^2(1)$. If $x \in S^2(1)$ then by (10) we get $\lambda = \varepsilon$, $\varepsilon = \pm 1$. The parametric equation of semi-Euclidean sphere in the adapted coordinates (u, φ) is:

$$\mathbf{r}(u, \varphi) = \varepsilon(\cosh \varphi, \sinh \varphi, u \exp \varphi), \quad (13)$$

where u is a basis coordinate, φ is a fiber coordinate. Different values of ε correspond to different connected components of semi-Euclidean sphere $S^2(1)$.

Let us define the action of the structure group S_1 on semi-Euclidean sphere. By (12) and the adapted coordinates of elements $a(0, \varepsilon_1, \psi)$, $x(u, \varepsilon, \varphi) \in S^2(1)$ we get:

$$u' = u, \quad \varepsilon' = \varepsilon \varepsilon_1, \quad \varphi' = \varphi + \psi.$$

This group also consists of two connected components.

The metric tensor for semi-Euclidean sphere has the matrix representation:

$$(g_{ij}) = \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}, \quad \text{rank}(g_{ij}) = 1$$

So the linear element of metric is:

$$ds_1^2 = -d\varphi^2. \quad (14)$$

3. SEMI-CONFORMAL MODEL OF THE SPHERE FIBRATION $(S^2(1), \pi, \mathbb{R})$

We consider the semi-conformal model of the fibration $(S^2(1), \pi, \mathbb{R})$. We project stereographically the sphere $S^2(1)$ from the point $N(1, 0, 0) \in S^2(1)$ (pole) to the equatorial plane $\mathbb{R}^2$ with equation $x_0 = 0$. We write the stereographic map in the coordinate form. For this we find the equation of line which pass through the pole and an arbitrary point of the sphere $X(x_0, x_1, x_2) : \tilde{x}_0 = 1 + (x_0 - 1)t, \tilde{x}_1 = x_1 t, \tilde{x}_2 = x_2 t$. When this line crosses the equatorial plane, we get the formulas of stereographic map $f : S^2(1) \rightarrow \mathbb{R}^2$ where $x_0 \neq 1$:

$$x = \frac{x_1}{1 - x_0}, \quad y = \frac{x_2}{1 - x_0}, \quad (15)$$

here $(x, y) \in \mathbb{R}^2$, (x_0, x_1, x_2) are coordinates on $S^2(1)$. An inverse map $f^{-1} : \mathbb{R}^2 \rightarrow S^2(1)$ when $x \neq \pm 1$ is:

$$x_0 = -\frac{1 + x^2}{1 - x^2}, \quad x_1 = \frac{2x}{1 - x^2}, \quad x_2 = \frac{2y}{1 - x^2}. \quad (16)$$

If we put formulas (15) into (13) then we obtain the relations between coordinates x, y and adapted coordinates u, φ which are on semi-Euclidean sphere:

$$f : \quad x = \frac{\sinh \varphi}{\varepsilon - \cosh \varphi}, \quad y = \frac{u \exp \varphi}{\varepsilon - \cosh \varphi}.$$

Then the inverse map is:

$$\varphi = \ln \left(\varepsilon \frac{x - 1}{x + 1} \right), \quad u = -\frac{2y}{(1 - x)^2}. \quad (17)$$

If we expand $\mathbb{R}^2$ to the semi-conformal plane C^2 by infinitely distant point and ideal line crossing this point, then the stereographic map f becomes diffeomorphic map $\tilde{f}$. This infinitely distant point is the image of point N . The ideal line is the image of straight line belonging to $S^2(1)$ and crossing the pole: $x_0 = 1, x_1 = 0$. Do not forget, that we need to exclude lines $x \neq \pm 1$ from the the plane $\mathbb{R}^2$.

Let us now consider the commutative diagram:

$$\begin{array}{ccc} S^2(1) & \xrightarrow{\tilde{f}} & C^2 \\ \pi \searrow & & \swarrow p \\ & \mathbb{R} & \end{array}$$

The map $p = \pi \circ \tilde{f}^{-1} : C^2 \rightarrow \mathbb{R}$ is defined by this diagram. We find the coordinate form of this map:

$$u = -\frac{2y}{(1 - x)^2}.$$

So, the map $p : C^2 \rightarrow \mathbb{R}$ defines the trivial principal fibration with the base $\mathbb{R}$ and the structure group S_1 .

Proposition. *The map $\tilde{f} : S^2(1) \rightarrow C^2$ is conformal.*

Proof. The metric on G induces the metric on C^2 . In the coordinates x, y it has the form:

$$d\tilde{s}^2 = -dx^2. \quad (18)$$

Let us find the metric of semi-Euclidean sphere from the metric on C^2 . By (17) we get $d\varphi = \frac{2}{x^2-1}$. So, by (14) and (18), we find:

$$ds_1^2 = \frac{4}{(x^2-1)^2} d\tilde{s}^2.$$

Hence, the linear element of semi-Euclidean sphere differs from the linear element of semi-plane by conformal factor and so, the map $\tilde{f}$ is conformal. $\square$

We find the fibers equations on C^2 . The 1-parametric fibers family of the fibration $(S^2(1), \pi, \mathbb{R})$ in the adaptive coordinates (13) is: $u = c$, $c \in \mathbb{R}$. By (17) we get the image of this family under map $\tilde{f}$:

$$y = -c/2 \cdot (x-1)^2. \quad (19)$$

The semi-conformal plane is also fibred by this 1-parametric family of curves. It is parabolas with axis $x = 1$ and top $(1, 0)$.

4. THE PROJECTIVE SEMI-CONFORMAL MODEL OF THE FIBRATION OF SPHERE $S^2(1)$

Now we construct the projective semi-conformal model of the sphere $S^2(1)$ and the principal fibration on it. We use normalization method of A.P.Norden [5]. A. P. Shirokov in his work [6] constructed conformal models of Non-Euclidean spaces with this method.

In a projective space P_n a hypersurface X_{n-1} as an absolute is called *normalized* if with every point $Q \in X_{n-1}$ is associated:

- 1) line P_I which has the point Q as the only intersection with the tangent space T_{n-1} ,
- 2) linear space P_{n-2} that belongs to T_{n-1} , but it does not contain the point Q .

We call them *normals of first and second types*, P_I and P_{II} .

In order that normalization be polar, P_I and P_{II} must be polar with respect to absolute X_{n-1} .

We expand the semi-Euclidean space ${}_2E_1^3$ to a projective space P^3 . Here ${}_kE_l^n$ denotes an n -dimensional semi-Euclidean space with metric tensor of rank k , l is the number of negative inertia index in a quadric form. We consider homogeneous coordinates $(y_0 : y_1 : y_2 : y_3)$ in P^3 , where $x_i = \frac{y_i}{y_3}$, $i = 0, 1, 2$. Thus $S^2(1) : x_0^2 - x_1^2 = 1$ describes the hyperquadric in P^3 :

$$y_0^2 - y_1^2 - y_3^2 = 0. \quad (20)$$

Here the projective basis (E_0, E_1, E_2, E_3) is chosen in the following way. The vertex E_0 of basis is inside the hyperquadric. The other vertices E_1, E_2, E_3 are on its polar plane, $y_0 = 0$. The line E_0E_3 cross the hyperquadric at poles $N(1 :$

The stereographic map of projective plane $P^2 : y_0 = 0$ to the hyperquadric (20) from the pole $N(1 : 0 : 0 : 1)$ is shown on the picture. Let $U(0 : y_1 : y_2 : y_3) \in P^2$. If $y_3 = 0$, then the line UN belongs to the tangent plane $T_N : y_0 - y_3 = 0$ of the hyperquadric (20) at the point N and in this case the intersection point of the line UN with the hyperquadric is not uniquely determined. If $y_3 \neq 0$, then the intersection point of line UN with the hyperquadric is unique. So, we choose the line $E_1E_2 : y_3 = 0$ as the line at infinity. In the area $y_3 \neq 0$ we consider the Cartesian coordinates $x_1 = \frac{y_1}{y_3}, x_2 = \frac{y_2}{y_3}$. Then the plane $\alpha : y_0 = 0, y_3 \neq 0$ becomes a plane with an affine structure A^2 . It is possible to introduce the structure of semi-Euclidean plane ${}_1E^2$ with the linear element

The hyperquadric and the plane α do not intersect or intersect in two imaginary parallel lines

The restriction of the stereographic projection to the plane α maps the point $U(0 : x_1 : x_2 : 1)$ into the point X_1

$$X_1(-1 - x_1^2 : 2x_1 : 2x_2 : 1 - x_1^2). \quad (23)$$

We construct an autopolar normalization of the hyperquadric. As a normal of first type we take lines with fixed center E_0 and as a normal of second type we

take their polar lines which belong to the plane α and cross the vertex E_2 of the hyperquadric. The line E_0X_1 intersects the plane α at the point

$$X(0 : 2x_1 : 2x_2 : 1 - x_1^2).$$

In this normalization the polar of the point X intersects the plane α on normal P_{II} . Thus for any point X in the plane α there corresponds a line which does not cross this point. It means that the plane α is also normalized. Normalization of α is defined by an absolute quadric (22).

We consider the derivative equations for this normalization. If we take normals of the first type with fixed center E_0 , then the derivative equations ([5], p.204) have the form:

$$\begin{aligned}\partial_i X &= Y_i + l_i X, \\ \nabla_j Y_i &= l_j Y_i + p_{ji} X.\end{aligned}\tag{24}$$

The points X, Y_i, E_0 define a family of projective frames. Here Y_i are generating points of normal P_{II} .

We can calculate the values $(X, X), (X, Y_i)$ on the plane α using the quadric form, which is in the left part of equation (20). So, $(X, X) = -(1 + x_1^2)^2$.

Let us find coordinates of metric tensor on plane α . So, we take the Weierstrass standartization

$$(\tilde{X}, \tilde{X}) = -1, \quad \tilde{X} = \frac{X}{1 + x_1^2}.$$

Then the coordinates of metric tensor are scalar product of partial derivatives $g_{ij} = -(\partial_i \tilde{X}, \partial_j \tilde{X})$:

$$(g_{ij}) = \begin{pmatrix} \frac{4}{(1+x_1^2)^2} & 0 \\ 0 & 0 \end{pmatrix}.$$

So, we get the conformal model of polar normalized plane $\alpha : y_0 = 0, y_3 \neq 0$ with a linear element

$$ds^2 = \frac{dx_1^2}{(1 + x_1^2)^2}.\tag{25}$$

It means that this non-Euclidean plane is conformal to semi-Euclidean plane ${}_1E^2$.

The points X and Y_i are polarly conjugated, $(X, Y_i) = 0$. From this equation and the derivative equations (24) we can get the non-zero connection coefficients:

$$\Gamma_{11}^1 = \Gamma_{12}^2 = \Gamma_{21}^2 = -\frac{2x_1}{1 + x_1^2}, \quad \Gamma_{11}^2 = \frac{2x_2}{1 + x_1^2}.$$

The sums $\Gamma_{ks}^s = \partial_k \ln \frac{c}{(1+x_1^2)^2}$ ($c = \text{const}$) are gradient, so the connection is equiaffine. Curvature tensor has the following form:

$$R_{121.}^2 = -R_{211.}^2 = -\frac{4}{(1 + x_1^2)^2}.$$

Ricci curvature tensor $R_{sk} = R_{isk}^i$ is symmetric: $R_{11} = \frac{4}{(1+x_1^2)^2}$. Metric g_{ij} and curvature R_{rsk}^i tensors are covariantly constant in this connection: $\nabla_k g_{ij} = 0$, $\nabla_l R_{rsk}^i = 0$.

Geodesic curves in this connection satisfy the following equations:

$$x_2 = A(x_1^2 - 1) + Bx_1 \text{ and } x_1 = 0,$$

where A and B are arbitrary constants. There are parabolas and lines orthogonal to axis Ox_1 .

Let us consider the fibration of this plane by double numbers subalgebra. We write the equations of fibers of semi-Euclidean sphere $S^2(1)$ in homogeneous coordinates:

$$\begin{cases} (y_0 - y_1)v - y_2 = 0, \\ y_0^2 - y_1^2 - y_3^2 = 0. \end{cases} \quad (26)$$

This 1-parametric family of curves fibers the hyperquadric and it defines a fibration on it. The stereographic projection of these fibers from the pole N to the plane α is:

$$x_2 = -v/2 \cdot (x_1 + 1)^2.$$

It is 1-parametric family of parabolas.

REMARK

We would obtain the similar results for the space of right cosets by Lie subgroup H_2 (it is the subgroup of invertible dual numbers) and the fibration of the group G by H_2 . However, H_2 is a normal divisor of the group G . Therefore, the spaces of right and left cosets coincide. We have the proposition:

Proposition *The fibration $(G, \pi', M' = G/H_2)$ determined by the formula*

$$\pi'(x) = \frac{x_1}{x_0} \quad (27)$$

is a trivial principal fibration over the real line $\mathbb{R} \setminus \{0\}$. The typical fiber is a plane without a line. The structure group is H_2 .

The equations of fibers are:

$$ux_0 - x_1 = 0, \quad u \in \mathbb{R}. \quad (28)$$

This is a 1-parametric family of planes with common axis Ox_2 . It is necessary to remove intersection points of these planes with planes: $x_0 \pm x_1 = 0$, $x_0 = 0$.

The left multiplications $x' = ax$ on the group G preserve fibres if and only if when $a \in H_2$.

The right multiplications: $x' = xb$ on group G forms the 3-parametric Lie group of linear transformations that preserve the fibration and induce a 1-parametric group of hyperbolic transformations on the base with two fixed points $u = \pm 1$:

$$u' = \frac{u + \alpha}{\alpha \cdot u + 1},$$

here $\alpha = \pi(b)$.

The right multiplications include involutions of hyperbolic type:

$$R(b) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ b_2 & b_2 & -1 \end{pmatrix}, R^2 = \text{Id}.$$

They induce the involution on the base:

$$u' = \frac{1}{u}.$$

The similar properties has the fibration of semi-Euclidean sphere by subgroup $S_2 = \{x \in H_2 | x_0^2 - x_1^2 = 1\}$ and its models.

ACKNOWLEDGEMENT

I would like to thank Professor Jiri Vanzura for fruitful discussions and support with writing this paper. This work was supported in part by grant No. 201/05/2707 of The Czech Science Foundation and by the Council of the Czech Government MSM 6198959214.

REFERENCES

- [1] B. N. Shapukov, Exercises on Lie groups and its applications, Moskow: RXD (2002).
- [2] Study E., Cartan E. Nombres complexes, Encyclopedie des sciences mathematiques pures et appliquees, t.1. Vol.1. (1908) 329-468.
- [3] V. V. Vishnevskii, A.P. Shirokov, V.V. Shurygin, Spaces over algebras, Izdat. Kazan. univ. (1985).
- [4] N. E. Belova, various articles and works published in Kazan, for example: Mat. vseros. molod. nauc. shkoli-konf. po mat. mod., geom. i algebre. Kazan.(1998) 169-174.
- [5] A. P. Norden, Spaces with affine connection, Moscow: Nauka (1976).
- [6] A. P. Shirokov, Non-Euclidean spaces, Izdat. Kazan. univ. (1997).
- [7] B. A. Rosenfeld, Geometry of Lie Groups, Kluwer (1991).